SPATIAL GEVENETRY OF THE HUMAN PELVIS

H. M. Reynolds 1

C. C. Snow 2

J. W. Young 3

Protection and Survival Laboratory Memorandum Reports are usually preliminary or progress-accounting documents intended for use within Government or by a specific non-Government activity, and as such, may not comply with DOT and FAA publications procedures and formats. Information centained in these reports is subject to additional evaluation or change on review of the data, conduct of additional testing or receipt of additional facts.

Protection and Survival Laboratory
Civil Aeromedical Institute
Pike Monroney Aeronautical Conter
Federal Aviation Administration
Oklahoma City, Oklahoma

1981

¹ College of Ostcopathic Medicine, Michigan State University

² Civil Aeromedical Institute, Federal Aviation Administration (Ret.)

³ Civil Aeromedical Institute, Federal Aviation Administration

INTRODUCTION

As a part of the human skeletal system, the pelvis is located at approximately the center of the body and forms a structure through which forces are transmitted. Under impact conditions, vertical forces are transmitted from the seat through the isohial tuberosities, and horizontal forces are transmitted from the seat belt, etc., through the ilium. When crash tests are conducted using anthropomorphic test devices to represent the human body, submarining of the test device occasionally occurs and has been attributed to the interaction of the test device pelvis geometry and the seat restraint system (Leung, et al., 1979). Thus, the geometry of the adult human pelvis is an important parameter in the dynamic behavior of the human body. The pelvis has also been used to determine sex of skeletal remains among disaster victims (Krogman, 1962). Differences in those dimensions which affect size and shape of the pelvic inlet and outlet (i.e., the birth canal) are used in this identification process. In addition, the obstetrical significance of the size and shape of the female pelvis has long teen noted and categorized (Cauldwell and Moloy, 1933).

This study, initiated to define and describe the spatial geometry of the adult pelvis, was jointly sponsored by the FAA (A':-IRS-CO), NHTSA (Interagency Agreement DOT-HS-6-01356), and AFOSAR (F-44620-7C-C-0015).

EXPERIMENTAL DECIGN

The features described for these various applications in the geometry of human skeletal anatomy have not been described in a systematic manner that would provide representative models of the complete human pelvis. As a result, an investigation has been conducted to measure the three-dimensional size and shape of the adult male and female pelvis. Specimens for this investigation were obtained from the Hamann-Todd skeletal collection in the Cleveland Museum of Natural History. The specimens were selected from nearly 3000 human skeletons of documented age, sex, and race. The skeletons came from the dissection room

population at Western Reserve School of Medicine between 1919 and 1939. A complete series of anthropometric dimensions were obtained by Dr. Tedd and his associates prior to the skeletonization of these cadavers. These data were used to select the specimens so that they represented as closely as possible the current general US non-institutionalized population. It was assumed that the size and shape of the pelvis would vary with respect to body size. That is, there would be a tendency (recognizing the importance of human variation in any correlative study) for large pelves to be associated with large body size and small pelves to associated with small tody size. Churchill and McConville (1976) have pointed out that body size can be represented by height and weight when attempting to match samples. As a result, specimens were selected from the Hamann-Todd skeletal collection on the basis of their height and weight with respect to the height/weight distribution in the Health and Examination Survey (NES) of 1961-1964 conducted by the US Public Health Service (Stoudt et al., 1965). Only 18-55 year old subjects in the HES raw anthropometric data were selected for analysis since it was reasoned they would be representative of the general US population of the middle to late 1970's. These data were available for reanalysis on the computer system at the University of Michigan.

The matching strategy was based upon the linear relationship between height and weight in the HES data. Figure 1 depicts the categories for both male and female body sizes.

Each category is defined by the intersection of regression lines which divide the theoretical ellipsoid representing the height/weight distribution into six categories for each sex.

Each category is coded so that they represent the following sizes:

- 1 Small stature/Heavy weight
- 2 Medium stature/Heavy weight
- 3 Tall stature/Heavy weight
- 4 Small stature/hight weight
- 5 Medium stature/Light weight
- 6 Tall stature/Light weight

These categories when combined into small (1 & 4), medium (2 & 5), and large (3 & 6) body sizes were established to divide the distribution into approximately 25 percent of the population at each of the extremes and 50 percent of the population in the middle (Table 1).




Figure 1. Sampling scheme for male and female specimens from the Hamann-Todd collection.

HEIGHT (om)

FEI-IALE

Sice Cell	HE	11—T	HES	H - T
**************************************	Ave. (3.0.)	Avc. (S.D.)	Avc. (S.D.)	Ave. (S.D.)
ı	164.6 (3.17)	165.1 (5.66)	152.6 (2.99)	151.5 (2.83)
2	173.5 (2.54)	172.9 (2.35)	160.2 (2.39)	160.4 (1.87)
3	182.0 (3.60)	182.8 (3.93)	168.7 (3.87)	169.1 (3.60)
4	164.9 (3.66)	164.1 (3.03)	152.5 (3.43)	153.9 (3.16)
5	173.9 (2.44)	174.4 (2.73)	160.9 (2.48)	161.5 (2.71)
6	182.5 (3.41)	184.8 (4.57)	168.9 (2.94)	170.7 (5.31)

MALE

WEIGHT (kg)

		MAJ-E	FEM	ALE
Size Cell	HES	11—T	HES	H-T
	Ave. (S.D.) Ave. (S.D.)	Ave. (S.D.)	Ave. (S.D.)
1	79.0 (9.00	79.7 (6.70)	71.7 (10.1)	74.8 (12.3)
2	85.9 (8.50) 63.1 (8.90)	77.3 (12.3)	73.8 (8.50)
3	92.7 (8.60) 91.5 (7.30)	82.3 (13.5)	78.7 (9;40)
4	62.1 (6.10) 63.3 (5.40)	52.4 (5.40)	46.9 (6.60)
5	67.6 (5.80	67.6 (6.80)	55.8 (5.40)	54.1 (6.70)
6	74.0 (6.00	/6.1 (8.20)	59.4 (5.40)	56.4 (9.30)

Table 2. A comparison of height and weight by size cell and sex (between HEC and Hamann-Todd specimens)

The corresponding sample sizes in the matched Marann-Todd specimens were selected to obtain subsamples large enough to represent each body size rather than to represent the distribution of body sizes in the general US population.

	HALE		F'E24ALJ	E
Size Cell	HES	H-T	HES	II-T
ı	291 (11%)	7	320 (10%)	16
2	658 (34%)	17	651 (20%)	17
3	339 (12%)	14	322 (10%)	11
4	361 (13%)	13	446 (14%)	12
5	761 (28%)	17	997 (31%)	19
6	344 (12%)	12	476 (15%)	10
Total	2754 (100%)	80	3212 (100%)	85

Table 1. Sample size in each size cell by sex

Selecting subjects from the Hamann-Todd skeletal collection was complicated by the fact that their weights were extremely low for their heights. As a result, a constant of 12.7 kg (28 lbs.) was added to the male weights and a constant of 14.1 kg (31 lbs.) was added to the female weights. The constants represent differences by sex between the average weight for the 18-70 year old males and females in the Health Examination Survey and the Hamann-Todd skeletal collection. Table 2 reports the average height and weight by sex and size cell for both the Health Examination Survey subjects as well as the Hamann-Todd cadaver population. The difference between heights are minimal, but even in the correction for weight there are substantial differences in weight. As previously pointed out, the size cells are defined primarily by height, and when the weight groups for the same heights are combined, the effect of weight is minimized on the categories defining body size.

EXPERIMENTAL METHODS

3-Dimensional Measuring

Following the reassembly of the pelvic ilia and sacral components into their correct anatomical position, a series of surface landmurks (Figures 3,4,5, and 6) for measurement procedures were placed on each specimen as defined in the listing of definitions (Appendix B). Specimens were first measured 2-dimensionally by conventional methods, then measured in the 3-dimensional format to establish the data base. Except for a few reference marks on the right hemipelvis, most of the landmarks were recorded on the left hemipelvis only. Assuming bilateral symmetry, the projection of left-registered points to the right side was accomplished by a computer program. The equipment used to register landmarks on the specimens consisted of a Hewlett-Fackard Model 9864A digitizer with input into a Hewlett-Packard Model 9820 A calculator. This system was modified by attaching a diagraph to the digitizing ourser of the plotting board so that the point of the diagraph needle would be directly vertical to the cross-hairs of the digitizer at all times. A potentiometer on the diagraph registers the height of the needle above the plotting board and provides Z - axis data to the computer. The specimen was suspended over the board in a position allowing access to all of the landmarks on the bonc. The diagraph is maneuvered so that the needle is in contact with a given landmark. A single signal, activated on the digitizer, registers the X, Y, and Z coordinates of the landmark in relation to an origin point on the board. The spatial location of each registered landmark can then be transformed in reference to an anatomical axis system which is based on a transverse plane formed by the midpoint of the antero-superior margin of the sacrum (promontorion) and the left and right anterior-superior iliao spines.

Repeated calibration runs on test specimens revealed a mean accuracy of ± 0.5 mm on computed point-to-point linear measurements. Such accuracy exceeds that obtained by conventional osteological techniques.




Figure 3. 3-Dimensional data points for polvis; Left side

Medial View






Figure 4. 3-Dimensional data points for pelvis: Left side





Posterior View

Anterior View

Superior View

Pigure 6. 3-Dimensional data points for sacrum

3-Dimensional Modeling

One objective of this study was to provide a series of full-scale, anatomically correct pelvic models for use in the design of anthropomorphic test devices. Master models for the small female, medium male, and large male, as defined by the three-dimensional data, were first sculptured in clay then duplicated in epoxy by a casting technique.

The modeling technique used a procedure similar to that in measuring the original specimens for three-dimensional data, except in reverse. From a series of five computer generated Y - Z coordinate point plots (unilateral format) for each pelvis size, a bilateral composite (mirror image) plot was made on transparent plastic and mounted on a horizontally stablized plate glass surface. A thin, rigid aluminum and plaster gauze framework was constructed for each model size to internally support and stabilize the irregular soft clay form. The framework was aligned and suspended above the composite Y - Z plot so that the orientation of the medel Y - Z plane would be parallel to that of the plot surface. The pelvis model Y - 7 plane, defined by the bilateral Anterior Superior Iliac Spine points (201) and the midsagittal Pubic Symphysis point (204), is the origin reference plane for all vertical X distance values. Using a diagraph instrument, the lower fixed pointer was aligned with a Y - Z coordinate plot point, then the upper movable pointer raised vertically to the appropriate X height value. The clay was extended and shaped to make surface contact with the upper pointer which defined the spatial location of that point. This procedure was repeated for all coordinate points to establish the threedimensional characteristics of the complete pelvis. Surface contouring between the coordinate point locations was accomplished on the basis of experienced judgement to be anatomically correct, but omitting fine detail which would serve no practical function.

The final epoxy models (Appendix C) were produced by the Ecanthropus Memorial Casting Laboratory, Santa Fe, New Mexico, which was also responsible for the design and fabrication of the casting molds for the master clay models. A dimensional accuracy of ± 2 millimeters for the final epoxy models is within typical limits of the combined measurement error of

anatomical specimens and variations in naturally occurring bilateral assymetry.

DATA ANALYSIS

Upon completion of the measurements at CAMI, the data were entered into the University of Michigan Computer System where all subsequent data analysis was performed. Since the data were obtained with each specimen in a rigid but controlled position with respect to the three-dimensional anthropometer and its three-dimensional axis system, a comparable anatomical axis system (Reynolds and Mubbard, 1980) was established for each specimen. The axis system is defined by three landmarks: right and left anterior superior iliao spines (201) and symphysion (204) (as shown in Figure 2).




Figure 2. Pelvic axis system depicting direction of axes and coordinate definitions

These three points define the Y - % plane in the following manner: construct a line passing from right to left anterior superior iliac spines to establish the +Y axis. Next establish a perpendicular to the Y axis passing through symphysion (204) in a -Z direction.

The intersection of the Y and Z axes establishes the origin of the axis system and a +X axis is defined normal to the Y - Z plane according to the right hand rule. All data are then transformed into the new axis system.

Data were checked for keypunch errors and erroneous values through use of X - VAL

(Kikta and Churchill, 1978). This editing program outputs extreme values for each variable
in addition to useful surmary statistics which allow the user to visually check the data
for outliers in each variable. Because of the reliability of the three-dimensional
authropometer, there have been no errors detected to date that can be attributed to electrical
read errors within the measurement system. There were a few keypunch errors detected and
corrections were made. A more serious error is obvious in the data which has been attributed to the axis system definition methodology. That is, when only three points in the
axis system definition exist, slight differences caused by observer error or variability in
the pelvis itself can produce a significant effect on the new data coordinates (Robbins,
1977). This source of variability has not been investigated to determine an optimum axis
system definition, and subsequent analyses may provide slightly different results than those
reported in this study.

Following completion of the data cdit, specimens were combined into the assigned size cells discussed previously. The grouped data for small (size cells 1 and 4), medium (size cells 2 and 5), and large (size cells 3 and 6) pelves were statistically summarized. These summary statistics (mean, standard deviation, and maximum - minimum values) were computed for the coordinate of each point. Since there were three (X, Y, Z) coordinates for each of the 123 points, a total of 369 variables were computed for the statistical summary. These data are reported in Appendix A.

In order to utilize these data for constructing the anatomical models, two-dimensional coordinate plots of the Y - Z plane were generated so that a perpendicular (X axis coordinate) defines a point above the plane for use by the sculpturer. There were five plots generated with three points in cormon on each plot: Promontorion (point 177),

heft Iliospinale, Summum (point 201), and Symphysion (point 204). These three points thereby provided a means of aligning each plot in the same relative position to construct a complete bilateral composite plot for sculpturing procedures.

CONCLUSION

In conclusion, these data will provide an estimate of the variation in pelvic geometry within the adult male and female civilian US population. The definition of axes systems using anatomical landmarks needs relatively "stable", palpable landmarks that are commensurately distant. This information can be obtained in a reasonably large scruple representative of extreme body sizes in the general US civilian population. Additional uses of these data will include specification of representative pelves for various sizes of anthropomerphic test devices, identification and description of cadavers in forensic investigations, and quantitative information on the shape and size variability of the adult human pelvis.

REFERENCES

- 1. Caldwell, W. E. and H. C. Moloy. 1933. Anatomical Variations in the Female Pelvis and their Effect in Labor with a Suggested Classification. Amer. J. Obs. & Gynec. 26:479-505.
- 2. Churchill, Edmund and John T. McConville. 1976. Sampling and Data Gathering Strategies for Future USAF Anthropometry. Final Report, AMRL-74-102, Wright-Patterson AFB, Ohio.
- 3. Kikta, Taul and Thomas Churchill. 1978. Editing Procedure for Anthropometry Data Survey. Technical Report, AMM-TR-78-38, Wright-Patterson AFB, Ohio.
- 4. Krogman, W. M. 1962. The Human Skeleton in Forensic Medicine. Charles C. Thomas, Springfield, Ill.
- Leung, Y. C., Tarrier, A. Gayou, P. Mairesse, A. Dolmas, and P. Banzet. 1979.
 A Comparison Between Part 572 Dummy and Human Subject in the Problem of Submarining.
 SAE \$791026. Proceedings of Twenty-Third Stapp Car Crash Conference. October 17-19, 1979. San Diego, California. pp. 675-719.
- 6. Reynolds, Herbert M. and Robert P. Hubbard. 1980. Anatomical Frames of Reference and Biomechanics. Human Factors 22 (2):171-176.
- 7. Stoudt, Howard W., Albert Damon, Poss McFarland, and Jean Roberts. 1965.
 Weight, Height, and Selected Pody Dimensions of Adults. National Center for
 Health Statistics, Series 11, Number 11. U.S. Dept. of Health, Education and
 Welfare, Public Health Service, Washington, D.C.
- 8. Robbins, P. H. 1977. Errors in Definition of an Anatomically-Rased Co-ordinate System Using Anthropometric Data. Fublished as a Task Report in A FOUNDATION FOR SYSTEMS ANTHROPOMETRY, Phase I, Interim Report to USAFOSR, Contract No. F 44620-76-C-0115, HSRI, The University of Michigan. (AD-A042890)

A PTORMAGA

COORD	LIATE		G - EDITATE FOLDT VALUES								
POL			MAL FRE	N1.81	ı	eriere	J.F.		LARGE MA	LE	
!!O	V/J2	1;	115711	Sib	11	MARIT	នរា	11	MEAN	SD	
177	х	27	- 56	1.1	32	- 59	9	26	- 57	10	
178	Y	27	0	3	32	o	5	26	0	4	
179	Z	27	18	15	32	12	16	26	16	8	
180	Ιx	27	- 6	1	33	9	1	26	- 1	7	
181	Y	28	- 5	2	33	- 4	2	26	- 5	2	
182	z	28	- 76	9	33	- 77	9	26	- 78	8	
183	x	28	6	10	33	- 9	1	26	2	7	
184	Y	28	5	2	33	4	2	26	5	2	
185	z	28	- 76	9	33	- 78	9	26	- 77		
186	x	28	- 87	9	33	- 96	9	26	-100	9 8	
187	Y	28	- 66	7	33	- 61	7	26	- 62	7	
188	Z	28	-102	14	33	-119	11	26	-120	9	
189	x	27	- 44	5	33	- 48	6	26	 50	6	
190	Y	28	- 76	25	33	- 82	5	26	- 86	5	
191	l z	28	- 47	66	33	- 66	7	26	- 67	7	
192	x	28	-120	11	33	- 134	9	26	-137	11	
193	Y	28	- 40	8	33	- 36	8	26	- 37	7	
194	2	27	34	17	33	31	16	26	35	13	
195	х	28	- 40	10	33	- 48	9	26	- 49	9	
196	Y	28	-114	11	33	-119	10	26	-125	8	
197	Z	28	53	Ö	.33	60	7	26	63	8	
198	x	28	0	0	33	0	0	26	0	0	
199	Υ	28	-109	12	33 .	-113	9	26	-115	8	
200	Z	23	0	0	33	0	0	18	0	0	
201	ķ	28	0	n	23	Õ	0	25	o	0	
202	ľΎ	28	100	12	33	117	8	26	115	8	
203	z	7	0	1)	33	0	o	15	0	o	
204	X.	27	Q	C	33	Q	0	26	0	o	
205	Y	21	0	0	53	0	0	14	0	0	
206	Z	28	- 76	ò	33	- 78	9	26	- 78	9	
207	x	28	- 63	12	33	- <i>(</i> Ú)	9	26	- 68	12	
208	Y	2 8	19	3	33	26	5	26	28	4	
209	z	28	31	15	33	29	15	26	33	8	
210	x	27	− (ð	10	32	- 83	ŝö	26	- 78	8	
211	Y	27	53	5	32	56	5	26	59	5	
212	z	27	21.	16	32	28	26	26	27	12	
213	x	25	- 79	10	32	- 87	10	24	- 90	10	
214	Y	25	38	5	32	44	5	24	46	7	
215	7.	25	48	15	32	48	13	24	53	8	
21g ·	X	28	- 74.	11	33	- 79	14	26	- 81	12	
217	Y	28	O	3	33	0	5	26	0	4	
218	2	28	37	14	33	33	14	26	39	8	
519	x	28	- 77	12	33	- 85	9	26	- 87		
220	Υ ,	28	13	4	37	17	5	26	15	13	
221	Z	28	47	15	33	41	15	26	38	6	
222	x	27	- 82	12	33	- 89	16	26	- 92	13	

C(K)RI)		COMPUTATE TOTAL VALUES								را. پاکانیشنسی،
1,011	rr:	:	MJ. FEV	VI-IR	1	adia: 13	J.E	<u> </u>	LAIGE MAI	LE
::0	AXIS	11	LEVI	SD	ij	MEAN	SD	11	MEAN	SD
223	Y	27	22	4	33	26	6	26	28	5
224	7.	27	52	15	33	48	18	26	56	8
225	X	28	- 83	11	33	- 97	41	26	- 93	11
226	Y	28	14	4	33	17	6	26	18	4
227	Z	28	39	15	33	38	20	26	42	9
228	Х	28	- 88	12	33	- 99	10	26	-100	11
229	Y	28	22	4	33	26	6	26	28	4
230	Z	28	44	19	33	40	14	26	45	9
231	X	26	- 79	12	21	- 81	31	26	- 87	12
232	Y	26	13	3	31	17	4	26	16	4
233	Z	26	35	14	31	31	18	26	38	8
234	х	24	- 85	14	31	- 94	15	23	- 95	13
235	Y	24	0	4	31	0	4	23	0	4
236	Z	24	27	15	31	24	15	23	28	8
237	X	25	- 96	13	32	-104	18	23	-107	13
238	Y	25	0	4	35	0	4	23	0	4
239	Z	26	38	10	35	32	15	23	40	9
240	x	20	-109	11	30	-122	10	21	-122	12
241	Y	20	0	4	30	0	4	21.	0	4
242	7.	20	36	19	30	35	16	21	40	11
243	х	25	-121	11	5.,	-134	9	20	-135	10
244	Y	25	0	3	2.1	0	4	20	0	4
2.45	Z	25	25	1.,	2:1	18	18	20	23	12
246	х	22	-127	10	20	-141	9	21	-145	8
247	Y	22	0	3	20	0	3	21	0	4
248	Z	22	10	19	20	- 2	2	21	3	12
249	X	2.1	-133	8	3.7	-139	6	25	-145	6
250	Y	24	0		30	0	3	25	0	4
251	::	24	-43	20	30	- 60	21	25	- 60	.17
252	x	28	-119	10	31	-130	8	25	-139	31
253	Y	28	28	5	31	31	5	24	30	8
254	7.	28	13	17	31.	4	13	25	9	40
255	X	27	-134	8	35	-143	7	26	-147	6
256	Y	27	0	3	32	0	4	26	0	3
257	Z	27	- 13	17	32	- 25	21	26	_ 24	18
258	X	STP	REF	P TTT	CTD	REF	POINT	STD	REF	POINT
259	Y	#	H	н	*	#	#	.	*	*
260	Z	*	99			4	*	•	**	*
261	X	27	- 76	8	32	- 82	8	26	- 82	7
262	Y	27	- 47	6	32	- 48	6	26	- 50	6
263	z	27	7	13	32	3	11	26	4	8
264	X	26	- 84	9	33	- 91	8	24	- 92	8
265	Ϋ́	26	- 49	6	3.3 2.2	- 43	6	24	- 49	4
266	z	26	19	1.:	33	17	12	24	20	9
267	x	26	- 77	10	32	- 84	8	25	- 85	9
268	Ÿ	26	- 53	5	32	- 53	6	25	- 56	6
			L							

COORD			COMPRIME FORT VALUES									
POI	T:	::	MALL FRO	ALE:	1	AN ENTER	14:	_	LARGE MA	ĿΣ		
1:0	ZIXA	и	HEIAH	SD	ון	PEAN	SD	N	MFAN	SD		
269	Z	26	32	13	32	31	12	25	35	10		
270	х	27	- 69	9	33	- 73	8	26	- 74	7		
271	Y	27	- 55	5	33	- 54	6	26	- 56	6		
272	Z	27	19	13	33	16	11	26	19	9		
273	х	26	- 02	ò	33	-100	8	26	-102	10		
274	Y	26	- 42	6	33	- 41	6	26	- 43	5		
275	Z	26	- 4	14	33	- 12	12	26	- 11	9		
276	x	25	-119	33	32	-123	8	25	-125	8		
277	Y	25	- 43	5	32	- 42	4	25	- 44	5		
278	2	25	1	24	32	- 12	14	25	- 11	11		
279	X	26	- 96	9	32	-106	8	25	-108	10		
280	Y	26	- 46	5	32	- 43	5	25	- 44	6		
281	7.	26	ŢIJ	14	3.3	4	12	25	7	11		
282	x	26	86	11	33	- 93	8	26	- 96	9		
283	Y	26	- 48	6	33	- 47	6	26	- 47	5		
284	Z	26	15	15	33	11	12	26	12	10		
285	х	28	– ფ	ò	33	- 75	8	26	- 75	9		
286	Y	28	40	6	33	51	5	26	47	20		
287	7.	28	2	13	33	- 5	11	26	- 9	7		
288	х	28	- 72	10	33	- 77	8	26	- 76	9		
289	Y	28	26	4	33	2.7	4	26	30	5		
290	7.	23	15	14	33	9	13	26	12	8		
291	X	27	- 69	12	33	- 74	9	26	- 73	10		
292	Y	27	ρ	3	35	o	4	26	0	6		
293	Z	27	17	1.1	33	ŗ	14	26	12	7		
294	x	28	- 83	13	32	- 69	10	26	- 87	10		
295	Ŷ	28	0	3	32	0	4	26	0,	3		
296	Z	28	11	15	35	3	13	26	6	7		
297	X	28	-105	12	33	-113	8	26	-111	9		
298	Ϋ́	28	0		33	0	3	26	0	3		
299	z	28	ì	16	33	- 9	14	26	- 7	9		
300	X	28	-113	10	33	-123	6	26	-121	8		
301	Ŷ	28	0	3	33	0	3	26	0	3		
302	z	28	- 4	19	23	- 16	15	26	- 13	11		
303	x	28	-118	10	33	-126	6	26	-127	7		
303	Ŷ	28	0	3	33	0	3	26	0	3		
305	ż	28	- 11	17	33	- 23	15	26	- 22	10		
306	X	26	-125	7	33	-132	5	26	-135	5		
307	Y	26	0		33	0	3	26	0	3		
308	Z	26	. – 28	15	3.3 2.3	- 10	17	26	- 38	12		
309	x	c	-124	6	18	-134	6	11	-137	5		
310	Y	9	0	2	18	0	3	11	0	4		
311	2	g	- 41	15	18	- 52	18	11	- 58	17		
312	x	22	-126	8	32	-1.72	7	26	-137	9		
313	Ϋ́	22	0	.4	32	0	4	26	0	ú		
314	Z	22	- 46	17	35	- 63	17	25	- 62	16		
		L	.,		<u>. </u>	-,,,	***					

Coopp.	HIATE:	1		C :	4-0117	V9'E : 1972	CHULLAN		•	
POI			ivil Pili			990108130			LARGE MA	
		r	I		•					1
::0	AXIS	1:	10001	::N	!!	i (Kari	SD	11	MEAN	SD
315	X	24	-12)	õ	32	-135	6	25	-140	6
316	Y	24	િ	3	32	9	5	25	9	4
317	Z	24	- 43	19	32	- 62	17	26	- 52	52
318	Х	22	-125	8	25	-132	6	24	-1.37	6
319	Y	22	27	3	25	29	9	24	30	6
320	Z	22	- 28	16	25	- 48	18	24	- 45	14
321	X	28	-105	9 (31	-118	7	26	-119	8
322	Y	28	42	4	31	44	10	26	44	6
323	Z	28	- · 5	14	31	- 18	14	26	- 19	12
324	X	STD "	RHF	HOINT !	CTD	RHF	POINT	STD	ref	POINT
325	Y	"		**	**	**	*	*	*	**
326	Z		"					•	*	*
327	X	28	9	3	33	- 3	3	26	0	3
328 329	Y Z	28	106	12	33	112	9	26	110	8
330		28	10	4 8	33	9	5	26	11	6
331	X Y	28 28	- 37	11	33	- 46	7 8	26	- 48	11
332	Z		122 38	8	33	133 42		26	134	7
333	X	28 28	- 15		33		10	26	46	9
334	Ϋ́	28	123	4 12	33	- 20	4 9	26	- 19	4
335	7.	28	?7	5	33 33	132 30		26	132	8
336	x	28	- 12	5	33	- 18	5 5	26	33	4
337	Ϋ́	28	114	12		123	9	26	- 18	5
337	7.	28	31	.12 .1	33	34		26	122	7
339	X	23	- 38	ម	35 33	- 47	5 9	26	38 49	4
340	Ϋ́	28	114	10	33		8	26		9
341	Z	28	54	9	?3	124 61	8	26	125 I 60	8
342	X	28	- 65	10	33	- 77	10	26 26	63	8
343	Ϋ́	28	- 65	i) 10	33	100	7	26	- 81 100	10
344	z	28	65	10	33	74	10	26	78	· 8
345	X	28	- 61	10	33	- 71	16	26	- 77	•
346	Ŷ	28	60 - 01	9	33	95	7	26	91	12 21
347	ż	28	66	10	33	73	10	26	76	16
348	x	28	- 76	13	33	- 88	12	26	- 91	13
349	Ϋ́	28	58	7	33	63	10	26	63	9
350	z	28	63	14	33	68	13	26	73	9
351	X	28	- 84	11	33	- 98	12	26	-103	12
352	Ϋ́	20	64	6	33	66	6	26	66	7
353	z	2ช	64	13	55	ଜ	13	26	73	8
354	x	28	- 70	12	33	- 03	11	26	– 97	12
355	Ŷ	28	58	7	33	ω ω	7	26	60	7
356	7.	28	63	13	3.7	63	12	26	72	8
357	x	27	· -1 09	10	33	-124	11	26	-130	12
358	Y	27	45	5	33	46	6	26	45	6
359	z	27	47	15	33	46	13	26	40	11
360	X	28	-101	11	33	-114	9	26	-117	12
									<u> </u>	J

C - 130					• • • •		TABLES.		-	
יוטו	rr:	:	::W. 1.35.		;	· 10115. • 4	11.6	<u> </u>	LANGE MA	LE
.,,	AVIS	•:	:8%::	;*0	::	HEAH	30	11	HEATH	SD
361	Y	28	38	5	33	38	5	26	38	6
362	Z	28	44	15	33	42	14	26	44	10
363	х	27	-119	11	33	-134	10	26	-139	10
364	Y	27	38	6	33	37	5	26	38	7
365	2	27	32	17	3 3 i	2 0 /	16	26	32	11
366	х	28	-118	9	33	-131	8	26	-135	8
367	Υ	28	35	4	33	35	4.	26	35	5
368	2	28	17	15	33	11	15	26	12	10
369	Х	28	-113	ö	33	-126	7	26	-130	8
370	Y	28	43	4	33	44	4	26	44	4
371	z	2 8	11	15	33	4	13	26	4	10
372	Х	28	-j10	ΰ	33	-123	10	26	-128	7
373	Υ	28	11	.3	33	45	4	26	46	4
374	2	28	- 6	15	33	- 11	13	26	10	10
375	х	28	- 97	7	33	-111	8	26	-113	7
376	Y	28	52	4	33	52	4	26	55	4
377	Z	28	- 7	14	33	- 17	12	26	- 17	8
378	х	28	- 84	Ö	33	- 97	7	26	-101	11
379	Y	28	26)	5	.33	58	4	26	57	15
380	Z	28	- .9	12	33	- 14	11	26	- 16	12
381	Х	28	- 74	5	33	- 85	5	26	- 88	5
382	Υ	28	60	5	33	66	4.	26	67	4
383	Z	28	- 19	10	33	- 21	9	26	- 20	7
384	х	28	- 78	5	33	- 80	4	26	- 91	5
385	Y	28	62	5	33	60	4	26	60	4
386	Z	28	- 30	10	33	- 45	9	26	~ 45	7
387	Х	CTD	REF.	I CTOT	Sib	EFF	THIE	STD	REF	POINT
388	Y	"	**	**	"	*	*	11	•	*
369	z	*	*	*	*	*	*	*	•	*
300	X	22	- 8월	6	28	- 90	32	22	-106	5
391	Y	22	48	24	56	54	23	22	48	5
392	3	22	- 55	22	28	- 56	21	22	∽ 65	8
303	X	27	- 6 5	6	31	- 96	5	26	-107	37
394	Y	27	60	5	31	54	4	26	55	5
395	Z	27	- 73	12	31	- 86	9	26	- 81	31
396	X	28	- 90	В	33	- 99	6	26	-104	8
397	Y	28	56		33	48	5	26	50	8
398	2	28	- 94	13	33	-111	10	26	-110	8
399	X	28	- 70	10	33	- 76	7	26	- 77	8
400	Y	28	43	5	53	39	5	26	41	7
401	7.	28	-114	11	33	-132	8	26	-136	9
402	X	28	- 73	9	53	- 81	7	26	- 82	8
403	Y	28	71	5	35	67	5	26	71	7
404	7	28	- 05	11	33	-114	9	26	-112	8
405	X	28	- 90	6	33	- 91	6	26	- 93	6
406	Y	28	74	5	33	72	4	26	74	6

Co ista				1;	! D1'#	GE COM	ANTORG	_	_	
POI	lit:	::	Mil Phi	AI.E	1	SPIDIUS: 1W	<u></u>		LARGE HA	LE
::0	SIXA	H	i ipati	នា	11	HEAH	នា	Ħ	MEAN	s
407	z	28	- 72	11	33	- 85	8	26	- 83	
408	X	28	- 87	ð	33	- 96	7	26	99	
409	Y	28	66	5	33	60	5	26	63	
410	Z	28	-100	12	33	-118	9	26	-120	
411	X	28	- 41	G	33	- 45	4	26	- 47	
412	Υ	28	33	3	33	29	3	26	30	
413	Z	28	- 95	8	33	-106	7	26	-109	
414	X	28	- 41	7	33	- 41	5	26	- 42	
415	Y	28	31	3	33	. 25	4	26	26	
416	Z	28	-107	9	33	-121	7	26	-123	
417	Х	28	- 48	6	33	- 53	5	26	- 53	ŀ
418	Y	28	53	3	33	50	3	26	52	
419	2	2ઇ	- 74	9	33	- 84	8	26	- 85	
420	X	28	- 64	7	33	- 69	5	26	- 71	
421	Y	28	50	4	33	46	4	26	47	
422	Z	28	- 94	10	33	-107	8	26	-110	
423	x	28	- 25	5	33	- 29	4	26	30	
424	Y	28	4	2	33	5	2	26	4	
425	Z	28	- 95	9	33	-104	8	26	-106	
426	Х	28	- 9	4	33	- 12	45	26	- 10	
427	Y	28	6	3	33	9	2	26	6	
428	Z	28	89	9	33	- 94	42	26	- 97	
429	l x	28	- 16	3	33	- 21	2	26	- 26	
430	Y	28	3	2	33	3	2	26	3	
431	z	28	- 81	G	33	- 85	8	26	- 81	
432	x	26	6	4	33	- 5	46	26	2	
433	ΙΥ	26	27	5	33	25	3	26	27	
434	7.	26	- 75	8	22	- 76	42	26	- 83	
435	l x	28	- 13	A	33	- 13	3	26	- 14	
436	Y	28	38	4	33	41	3	26	41	
437	z	28	- 71	8	33	- 78	7	26	- 79	
438	х	28	- 11	ŝ	32	- 13	3	26	- 12	
439	Y	28	24	4	38	33	3	26	34	
440	Z	28	- 64	łş	32	- 69	7	26	- 67	
441	х	28	- 27	4	33	- 2)	3	26	- 29	
442	Y	28	33	3	33	34	3	26	34	
443	2	28	~ 73	8	33 -	- 79	8	26	- 79	
444	х	28	- 21	4	33	- 21	4	26	- 22	1
445	· · · Y	28	63	é	.33 \$	64	5	26	59	"
446	Z	28	- 46	8	33	- 53	7	26	- 66	6
447	x	STD	KEF	10IHr	STD	ref	POINT	STD	RFF	PC
448	Y	H	*	*	""	,,,,	W	*	n	١ '`
449	z	#	*	**			**	*		
450	x	28	- 53	5	33	· - 58	5	26	- 60	נו
451	Y	28	62	6	33	61	5	26	- 6 54	3
· 	,z		~~	· ·	. ~~		_	. ~~	J 7	ı

COO!U	HATE		COMBINATE FOLIT VALUES								
POIN	rr:		MALL FIM	AI-E	1	SDIM IV	J.E		LARGE MA	LE	
::0	AYIS	1;	i:A:B:	cn	11	HEAH	SD	11	MEAN	SD	
453	x	28	- 35	-1	33	- 39	4	26	- 44	18	
454	Y	28	52	5	33	51	4	26	44	31	
455	Z	28	- 48	8	33	52	7	26	- 55	23	
456	х	28	- 22	3	33	- 24	2	26	- 27	9	
457	Y	28	31	3	33	32	3	26	26	28	
458	2	28	- 68	B	33	- 73	7	26	- 75	21.	
459	х	28	- 79	7	33	- 87	4	26	- 90	11	
460	Y	28	56	5	33	50	4	26	45	31.	
461	Z	28	- 77	11	33	- 90	.9	26	- 91	18	
462	x	28	− 6 5	5	33	- 76	4	26	79	10	
463	Y	28	57	4	33	53	3	26	48	32	
464	$\overline{\mathbf{z}}$	28	- 58	10	33	- 68	9	26	- 70	19	
465	x	28	- 60	5	33	- 73	4	26	- 68	11	
466	Ŷ	28	61	5	33	73 59	4	26	52	33	
467	z	28	- 38	10	33	- 42	26	26	- 49	22	
468	x	28	- 49	6	33	- 56	5	26			
469		28	77				6		- 59	14	
409 470	Y	28 28	- ''	ા 9	33	80		26	72	37	
	Z				33	- 5	8	26	- 9	3	
471	X	28	- 46	7	33	- 55	6	26	- 64	52	
472	Y	28	90	9	33	96	7	26	87	41	
473	Z	28	17	8	33	15	8	26	12	35	
474	х	28	- 42	7	33	- 51	7	26	 57	19	
475	Y	28	101	10	33	109	· 8	26	99	44	
476	2	28	36	9	33	37 .	7	26	32	38	
477	х	28	- 62	8	33	- 69	9	26	- 74	16	
478	Y	2 8	75	6	33	82	5	26	74	38	
479	Z	28 1	22	11	33	17	9	26	12	32	
480	х	28	- 41	6	33	- 40	4 .	26	~ 53	17	
481	Y	28	90	8	33	96	6	26	86	· 41.	
482	Z	28	13	8	33	10	6	26	4	32	
483	X .	28	- 21	4	33	- 24	3	26	- 28	16	
484	Y	28	'ဝှဝ	9	33	105	6	26	93	43	
485	Z	28	7	4	33	5	4	25	- 2	34	
486	x	28	- 12	3	33	- 15	3	26	- 19	15	
487	Y	28	98	8	33	104	6	26	95	43	
488	Z	28	 9	4	33	- 12	4	26	- 18	30	
489	X,	28	·- 14	5	33	- 16	5	26	- 18	13	
490	Y	28	90	6	33	95	4 ,	26	87	42	
491	2	28	- 27	6	33	~ 33	6	26	- 36	27	
492	х	28	- 24	5	33	- 25	4	26	- 29	12	
493	Y	28	84	5	33	87	5	26	80	40	
494	z	28	39	6	33	- 45	7.	26	- 49	24	
495	x	STD	REF	TUIDIT	STD	REF	POINT	STD	REF	POINT	
496	Ŷ	(17.7)	No.P	1	810	M ·	POMIT	21.0	NEP M	FOTIAL	
497	z	*		"		H		<u>.</u>			
498	X	28	- 35		33	- 40					
720	_ ^	40	_ 55	4	33	- 40	3	26	- 45	15	

COORD	CIATE	C EDINATE FORTE ANDRES								
POI	VT:	::	MIN FRO	/14°	1	NI SUIGE	e de la companya de l		LAIRGE MA	LE
110	axis.	Ħ	1:A:A:1	::0	11	ITEMI	SD.	31	MEAN	SD
499	Y	28	101	8	33	108	6.	26	99	44
500	Z	28	4	5	33	1	4	26	- 5	31
501	X	28	- 63	6	33	- 70	5	26	- 86	14
502	Y	28	78	6	33	89	5	26	77	38
503	Z	28	10	8	33	5	7	26	3	30
504	X	28	- 95	8	33	-103	14	26	-112	14
505	Y	-28	60	5	33	63	4	26	56	34
506	2	28	16	13	33	7	12	25	5	29
507	X	28	- 45	6	33	- 57	7	26	- 62	17
508	Y	28	102	9	33	110	7	26	101	45
509	Z	28	26	9	33	27	6	26	22	36
510	X	28	- 50	5	33	- 60	5	26	- 65	14
511	Y	28	90	8	33	95	6	26	88	41
512	Z	28	1	9	33	- 1	6	26	~ 5	30
513	X	28	- 60	5	33	- 71	4	26	- 84	56
514	Y	28	84	7	33	87	5	26	81	38
515	Z	28	- 24	ò	33	30	7	26	- 28	8
516	X	28	- 72	5	33	- 84	5	26	- 96	53
517	Y	28	80	6	35	81	5	26	76	36
518	2	28	- 48	10	33	- 58	8	26	- 54	9
519	Х	28	- 24	5	33	- 24	4	26	- 31	23
520	Y	28	66	4	33	68	4	26	62	36
521	Z	28	- 63	8	33	- 71	7	26	- 75	21
522	Х	28	- 31	5	33	- 34	5	26	~ 39	11
523	Y	23	0]	5	33	96	5	26	90	42
524	Z	28	- 40	7	33	- 44	6	26	- 48	24
525	Х	28	- 63	6	3.5	- 73	5	26	- 77	10
526	Y	28	90	5	33	94	6	26	87	41
527	Z	28	- 51	8	33	- 57	8	26	- 60	20
528	Х	28	- 54	6	33	- 61	5	26	- 64	10
529	Y	28	71	5	33	73	5	26	67	37
530	Z	28	- 7 7	បំ	33	– 88	7	26	- 90	18
531	XI	28	- 51	4	33	- 56	4	26	- 60	11
532	Y,	28	62	5	33	62	6	26	56	34
533	Z	28	- 44	9	33	- 50	8	26	- 53	22
534	X	27	- 43	5	33	- 48	5	26	- 52	10
535	Y	27	61	4	3.	83	5	26	78	39
536	2	27	- ,50	7	33	- 65	7	26	- 69	20
537	X	27	- 76	7	32	- 82	9	26	- 81	7
538	Y	27	- 47	6	32	- 45	6	26	- 46	6
539	7.	27	- 1	13	32	- 8	11	26	- 6	8
540	Х	28	- 51	5	33	- 55	G	26	- 58	13
541	Y .	28	63	6	33	63	5	26	55	34
542	Z	28	- 19	9	33	- 24	9	26	- 28	26
543	X	28	- 47	6	33	- 4)	5	26	- 51.	5
544	Y	28	28	4	33	21	4	26	21	4
545	Z	28	<u> </u>	9	33	-114	8	_ 26	-115	8

Definitions of Pelvic Landmarks

Each descriptive surface landmark (measuring point) used in this study is identified in three-dimensional space as a set of three sequential numbers to represent the X, Y, and Z coordinate values. The first number in each set refers to the X coordinate value and is always used as the identifying reference number for each measurement point. These sequential number sets appear with each of the following measurement point definitions.

<u>Promontorion 177-179</u>: The mid-point of the anterior-superior margin of the first sacral segment (excluding exotoses).

Right Superior Pole, Public Symphysis 180-182: The point at the intersection of the longitudinal mid-line axis of the public symphysis with the superior margin of the symphyseal face of the right innominate bone.

Left Superior Pole, Fublic Symphysis 193-185: The point at the intersection of the longitudinal mid-line axis of the public symphysis with the superior margin of the symphyseal face of the left innominate bone.

Ischiale 186-188: The immominate rests on its medial surface with the iliac blade and public symphysis in contact with the horizontal surface of an osteometric board. Hove the bone into the right angle corner of the board in such a way that the superior border of the iliac crest is in contact with one of the vertical plates and the anterior forder of the iliac crest and the public bone are in contact with the second vertical plate of the osteometric board. Ischiale is the highest point on the ischial tuberosity when the innominate bone is in the correct position.

<u>H-Point 189-191</u>: A plexiglass hemisphere is selected which best fits the acetabulum of the left innominate. The hemisphere is positioned so that the anterior extremity of one of the perpendicular diameter lines is opposite Point 519 (Acetabulion Anterior). H-Point is the center point of the hemisphere surface.

Iliocristale, Posterior 192-194: The innerinate rests on its medial surface with the iliae blade and public symphysis in contact with the horizontal surface of an esteometric board. Here the bene into the right angle corner of the board in such a way that the superior border of the iliae crest is in contact with one of the vertical plates and the anterior border of the iliae crest and the public bone are in contact with the second vertical plate of the esteometric board. Iliocristale Posterior, the posterior superior iliae spine, is defined as the point along the posterior border of the iliae crest in centact with the vertical plate of the esteometric board. In cases where a large area is in contact with the board, the landmark is taken as the mid-point of the centact area.

Iliocristale, Summm 195-197: The innominate rests on its medial surface with the iliac blade and pubic symphysis in contact with the horizontal surface of an osteometric board. The bone is moved into the right angle corner of the board in such a way that the superior torder of the iliac crest is in contact with one of the vertical plates and the anterior border of theiliac crest and the pubic bone

are in contact with the record vertical plate of the ostcometric board. Iliocristale Summum is defined as the point along the superior border of the iliac crest that is in contact with a movable vertical plate oriented at right angles to the vertical plates of the osctometric board. In cases where a large area is in contact with the board, the landmark is taken as the mid-point of the contact area.

Right Iliospinale, Summum 198-200: The right innominate rests on its medial surface with the iliac blade and pubic symphysis in contact with the horizontal surface of an esteemetric board. Here the bone into the right angle corner of the board in such a way that the superior border of the iliac crest is in contact with one of the vertical plates and the anterior border of the iliac crest and the pubic bone are in contact with the second vertical plate of the esteemetric board. Iliospinale Summum, the anterior superior iliac spine, is defined as the point along the anterior border of the iliac crest in contact with the vertical plate. In cases where a large area is in contact with the board, the landmark is taken as the mid-point of the contact area.

beft Iliospinale, Summur 201-203: The left innominate rests on its medial surface with the iliac blade and pubic symphysis in contact with the horizontal surface of an esteometric tourd. Here the lone into the right corner of the board in such a way that the superior lorder of the iliac crest is in contact with one of the vertical plates and the anterior border of the iliac crest and the pubic bone are in contact with the record vertical plate of the esteometric board. Iliospinale Summum, the anterior superior iliac spine, is defined as the point along the anterior torder of the iliac crest in contact with the vertical plate. In cases where a large area is in contact with the board, the landmark is taken as the mid-point of the contact area.

Pubic Symphysis 204-206: This point is the mid-point of a line between point 180, Right Superior Pole, Fulic Symphysis, and point 183, Left Superior Pole, Pubic Symphysis. It is not a landmark that is measured, but a point in space that is mathematically constructed.

Lateral Point on First Sacral Vertebral Pody 207-209: The most lateral point on the left articular surface of the first sacral body. In cases with "lipping" present, the point was estimated as the most lateral point on the superior surface that would be found in the general contour that would have been found without any "lipping".

Lateral Alax-Auricular Point 210-212: The point is found on the superior aspect as the most lateral point along the superior margin of the articular surface.

Posterior Alar-Auricular Point 213-215: The point is found on the superior aspect as the most postero-lateral boney projection on the wing of the sacrum.

Posterior Point on First Sacral Vertebral Fody 216-218: The Mid-point of the postero-superior margin of the first sacral segment. This point is posterior to Promontorion, point 177.

Superior Articular Facet: Medial Superior 219-221: The most superior point on the medial side of the superior articular facet of the sacrum.

Superior Articular Facet: Lateral Superior 222-224: The most superior point on the lateral side of the superior articular facet of the sacrum.

Superior Articular Facet: Medial Inferior 225-227: The most inferior point on the medial side of the superior articular facet of the sacrum.

Superior Articular Facet: Lateral Inferior 228-230: The most inferior point on the lateral side of the superior articular facet of the sacrum.

Sacral Ganal, Anterior Wall 231-233: The most lateral point on the lateral wall of the sacral canal. This point is found when the sacrum is oriented such that Point 177, Promontroion, Point 207, Lateral Point on First Sacral Vertebral Body, and Point 216, Posterior Point on First Sacral Vertebral Body, form a horizontal plane. The sacrum is viewed in the posterior aspect.

Sacral Canal, Anterior Floor 234-236: The sacrum is viewed in the posterior aspect while the sacrum is oriented in a horizontal plane defined by Point 177, Promontorion, Point 207, Lateral Point on First Sacral Vertebral Body, and Point 216, Posterior Point on First Sacral Vertebral Body. The landmark is defined as a point on the posterior body of the first sacral vertebral body at the intersection of a horizontal plane tangent to the anterior roof point of the sacral canal (Point 237) and a perpendicular line passing through the Posterior point on the First Sacral Vertebral Body Point 216.

Sacral Canal, Anterior Roof 237-239: The most anterior-superior point of the roof of the sacral canal.

Dorsal Spine of the First Sacral Vertebrae 240-242: The most posterior point on the dorsal spine of the first sacral segment. In the instance of a continuous boney ridge, the point is estimated by reference to the superior and inferior margins of the laminar plates that join to form the dorsal spines.

Dorsal Spine of the Second Sacral Verteirae 2-13-245: The most posterior point on the dorsal spine of the second sacral segment. In the instance of a continuous boney ridge, the point is estimated by reference to the superior and inferior margins of the laminar plates that join to form the dorsal spines.

Dorsal Spine of the Third Sacral Vertebrae 2:6-248: The most posterior point on the dorsal spine of the third sacral segment. In the instance of a continuous boney ridge, the point is estimated by reference to the superior and inferior margins of the laminar plates that join to form the dorsal spines.

Caudion, Posterior 249-251: The mid-point of the posterior-inferior margin of the last sacral segment. Herphological observations of the sacral segments should be examined to determine the exact sacral or coccygeal vertebrae upon which Point Caudion is located.

Sacral Canal, Posterior Roof 755-257: The point is located in the mid-sagittal plane at the most inferior aspect of the sacral canal roof.

Posterior Sacral Tubercle 252-254: The sacrum is viewed from the lateral aspect while oriented in a horizontal plane defined by Point 177, Promontorion, Point 207, Lateral Point on First Sacral Vertebral Body, and Point 216, Posterior Point on First Sacral Vertebral Body. The landmark is defined as the most posterior point on the posterior sacral tubercle adjacent to the 2nd vertebral foramen.

Sacroiliac Mid-Point 261-263: The point that lies at the intersect of the lines which bisect the superior and inferior poles of the sacro-iliac joint.

Superior Lole, Inferior Hargin Mid-Point 26:-266: A point along the inferior margin of the superior pole of the sacre-iliac joint surface that lies on a perpendicular line that bisects the line passing between Point 261, Sacre-iliac Mid-Point and Point 267, Superior Pole.

Superior Pole 267-269: A point on the posterior margin of the sacro-iliac joint surface that lies on a line bisecting the superior pole of the joint surface.

Superior Late, Superior Margin Mid-Foint 270-272: A point along the superior margin of the superior pole of the sacro-iliac joint surface that lies on a perpendicular line that bisects the line passing between Point 261, Sacro-iliac Mid-Point and Point 267, Superior Pole.

Inferior Lobe, Anterior Margin 273-275: A point along the anterior margin of the inferior lobe of the sacro-iliac joint surface that lies on a perpendicular line bisecting the line between Point 261, Sacro-iliac Mid-Point and Point 276, Inferior Pole.

Inferior Pole 276-278: A point on the inferior margin which lies on a line bisecting the inferior pole of the sacro-iliac joint

Inferior Lobe, Posterior Margin 279-291: A point along the posterior margin of the inferior lobe that lies on a perpendicular line bisecting the line between Point 261, Sacro-iliac Mid-Point and Point 276, Inferior Pole.

<u>Posterior Angle 282-284:</u> The most autorior point along the posterior margin of the sacro-iliac joint.

Anterior Alar Point 28-287: This point is found on the lateral aspect of the sacrum and is defined as the mort anterior point on the anterior margin of the articular surface of the sacro-iliae joint.

Hid-Alar Point 288-2.0: This point is defined on the anterior aspect of the sacrum as the mid-point of a line on the anterior surface of the alar wing between Point 201, Center of First Sacral Pertobeal Body and Point 285, Anterior Alar Point.

Center of First Sacral Vertebral Body 291-293: This point lies on the base of the sacrum equidistant between Point 177, Promontorion and Point 216, Posterior Point on the First Sacral Body.

First Sacral Segment Union Foint 294-296: This point lies in the mid-line of the sacrum defined as a line parting equidistant between segment sacral spertures (medial edge of vertebral canals). The first union point is located at the intersection of the vertical mid-line axis and the anatomical union of the first and second sacral vertebrae. If this closure is incomplete, the point is reconstructed in clay mid-way between adjacent vertebrae. This point is based on anatomical, not topographical, observations and is an estimate of osteological closure lines. The point may not be located at the most anterior aspect of this union area.

Second Sacral Segment Union Point 297-299: This point lies in the mid-line of the sacrum defined as a line passing equidistant between segment sacral apertures (medial edge of vertebral conals). The record union point is located at the intersection of the vertical mid-line axis and the anatomical union of the second and third sacral vertebrae. If this closure is incomplete, the point is reconstructed in clay mid-way between adjacent vertebrae. This point is based on anatomical, not topographical, observations and is an estimate of esteological closure lines. The point may not be located at the most anterior aspect of this union area.

Inflexion 300-302: This point is located at the intersection of the longest perpendiculars from the two extreme anterior ends of the sacrum. Inflexion is a geometric point which uses either L5 or Sl at the proximal end and Caudion at the distal end.

Third Sacral Segment Union Point 303-205: This point lies in the mid-line of the sacrum defined as a line passing equidistant between segment sacral apertures (medial edge of vertebral canals). The third union point is located at the intersection of the vertical mid-line axis and the anatomical union of the third and fourth sacral vertebrae. If this closure is incomplete, the point is reconstructed in clay mid-way between adjacent vertebrae. This point is based on anatomical, not topographical, observations and is an estimate of osteological closure lines. The point may not be located at the most anterior aspect of this union area.

Fourth Sacral Segment Union Point 206-308: This point lies in the mid-line of the sacrum defined as a line parsing equidistant between segment sacral apertures (medial edge of vertebral canals). The fourth union point is located at the intersection of the vertical mid-line axis and the anatomical union of the fourth and fifth sacral vertebrae. If this closure is incomplete, the point is reconstructed in clay mid-way between adjacent vertebrae. This point is based on anatomical, net topological, observations and is an estimate of osteological closure lines. The point may not be located at the most anterior aspect of this union area.

Fifth Sacral Segment Union Point 307-311: This point lies in the mid-line of the sacrum defined as a line passing equidistant between segment sacral apertures (medial edge of vertebral canals). The fifth union point is located at the intersection of the vertical mid-line axis and the anatomical union of the fifth and sixth sacral vertebrae. If this closure is incomplete, the point is reconstructed in clay mid-way between adjacent vertebrae. This point is based on

anatomical, not topographical, observations and is an estimate of osteological closure lines. The point may not be located at the most anterior aspect of this union area.

Caudion, Anterior 312-314: This point is at the mid-point of the antero-inferior margin of the last sacral segment.

Inferior Sacral Angle 318-320: This point is located, viewed anteriorly, at the intersection of lines approximating the latero-inferior aspect of the sacral body and the inferior margin of the sacral body.

Inferior Sacro-Iliac Junction 321-323: A point on the anterior edge of the sacro-iliac joint where the inferior curve of the sacral articular margin intersects the anterior sacral surface.

Iliospinale, Cristale 327-329: This point is observed in a posterior-anterior view across the iliac blade. It is defined as the most medial point on the medial margin of the anterior portion of the iliac crest.

Iliocristale, Anterior Lateral 330-332: The most lateral point on the lateral border of the iliac crest in the iliac pillar.

Anterior Segment Mid-Point, Lateral 333-335: A point that is located at the intersection of the lateral iliac crest edge and a line bisecting and perpendicular to a line passing through Point 339, Iliocristale Summum, and Point 201, Right Iliospinale Surrum.

Anterior Segment Mid-Loint, Medial 336-338: A point that is located at the intersection of the medial iliac crost edge and a line bisecting and perpendicular to a line passing through boint 337, Iliocristale Summum, and Point 201, Right Iliospinale Summum.

Iliocristale, Summum 337-3:1: The right innominate rests on its medial surface with the iliac blade and pubic symphysis in contact with the horizontal surface of an osteometric board. The innominate is neved into the right angle corner of the board in such a way that the superior border of the iliac crest is in contact with one of the vertical plates and the anterior border of the iliac crest and the pubic bone are in contact with the second vertical plate of the osteometric board. Iliocristale Summum is defined as the point along the superior border of the iliac crest in contact with the vertical plate. In instances where a large area is in contact with the plate, the landmark is taken as the mid-point of the contact area.

Posterior Segment Point, Lateral 1 342-344: Petween Point 339, Iliooristale Summum, and Point 366, Posterier Superior Iliospinale, on the iliao crest surface, three equidistant marks are made as perpendicular projections intersecting the medial and lateral margins. Lateral 1 point is located at the intersection of the most anterior mark and the lateral crest margin.

Posterior Segment Feint, Medial 1 345-347: Between Point 339, Iliocristale Summum, and Point 366, Fosterier Superior Iliospinale, on the iliac crest surface, three equidistant marks are made as perpendicular projections intersecting the

Posterior Iliac Notch Point 369-272; The deepest point on the posterior iliac border between Point 366, Posterior Superior Iliospinale, and Point 372, Posterior Inferior Iliospinale.

Posterior Inferior Iliospinale 372-374: The most projecting point on the posterior auricular margin.

Bouisson Tubercle 375-380: The most preminent point on the tubercle of Bouisson located at the apex of a tubercle formed by the origin of m. piriformis.

Superior Sciatic Notch Point 378-380: The point equidistant between Point 381, Apex of Sciatic Notch, and Point 375, Bouisson Tubercle. It is on the superior margin of the sciatic notch.

Apex of Sciatic Notch 381-383: The point on the sciatic notch border at the greatest perpendicular distance from a projected line between Point 372, Posterior Inferior Iliospinale, and Point 390, Ischial Spinale.

Anterior Sciatic Notch Point 384-386: The point on the inferior margin of the sciatic notch mid-way between Point 390, Ischial Spinale, and Point 381, Apex of the Sciatic Notch.

Ischio-Spinale 390-392: The superior-rosterior angle of the ischial spine.

Posterior Ischial Border Point 393-395: The point on the posterior margin of the ischial border equidistant between Point 390, Ischio-Spinale, and Point 396, Medial Tuberosity Point.

Hedial Tuberosity Point 306-398: The most medial point on the medial margin of the ischial tuberesity when the innominate is oriented in the anatomical position.

Inferior Tuberosity Point 399-401: The point of convergence of the medial and lateral margins of the ischial tuberosity.

<u>Lateral Tuberosity Point 402-404</u>: The most lateral point on the lateral margin of the isohial tuberosity with the immominate held in the anatomical position.

Superior Tuberosity Point 405-407: The most superior point on the margin of the ischial tuberosity with the innominate held in the anatomical position and viewed from the lateral aspect.

Ischiale 408-410: The right innominate rests on its medial surface with the iliac blade and pubic symphysis in contact with the horizontal surface of an osteometric board. The bone is moved into the right angle corner of the board in such a way that the superior border of the iliac crest is in contact with one of the vertical plates and the anterior border of the iliac crest and the pubic bone are in contact with the second vertical plate of the esteometric board. Ischiale is defined as the highest point on the ischial tuberosity from the surface of the esteometric loard.

Superior Ischial Public Ramus 411-413: The point equidistant between Point 180, Right Superior Pole, Public Symphysis, and Point 408, Ischiale, on the superior margin of the ischial public ramus.

Anterior Inferior Ischial Pubic Farus 414-416: Apoint on the anterior inferior margin of the ischial pubic ramus passing through Point 411, Superior Ischial Pubic Ramus, at right angles to the line joining Point 180, Right Superior Pole, Pubic Symphysis, and Foint 408, Ischiale.

Obturator Tubercle Point 417-419: The point on the lateral margin of the obturator foramen generally marked by a small tubercle. In instances where there are two tubercles, the apex of the most superior tubercle is marked as the point with the innominate oriented in the anatomical position.

Inferior Obturator Foramen Point 420-422: The point at which the greatest ourvature occurs on the posterior-inferior margin of the obturator foramen.

Inferior Symphyseal Pole 423-4.5: This point is located at the inferior intersection of the anterior and posterior margins of the symphyseal surface.

Anterior Symphyseal loint 426-428: This point is located on the anterior margin of the pubic symphysis where it is intersected by a line perpendicular to the longitudinal axis equidistant between the polar Points 180 and 423.

Posterior Symphyseal Point 429-431: This point is located on the anterior margin of the pubic symphysis where it is intersected by a line perpendicular to the longitudinal axis equidistant between polar Points 180 and 423.

432-434

Publisherale: This point is located at the anterior most projecting point of the summit of the public tuberale with the innominate oriented in the anatomical position.

Anterior Public Ramus Foint 435-437: The point on the anterior surface of the public ramus surking the intersection of a line passing through Point 456, Anterior Inlet Point, at right angles to the line of the pelvic inlet and the line extending from Point 432, Pubotubercle, to Point 519, Acetabulion, Anterior.

Superior Pubic Riemus Point 438-440: A point on the margin of the pubic crest crossed by a line passing through Point 456, Anterior Inlet Point, at right angles to the line of the pelvic inlet.

Inferior Public Partus Leint 441-443: The point on the superior border of the obturator foramen crossed by a line passing through Point 456, Anterior Inlet Point, at right angles to the pelvic inlet line.

Public Eminence Point 444-440: The surmit of the ilio-pectineal emminence.

Posterior Inlet Foint 450-452: The posterior most point subdividing a straight line into four equidist and segments from Foint 285, Anterior Alar Point, to Foint 429, Fosterior Symphysical Foint.

Intermediate Inlet Point 453-455: The intermediate most point subdividing a straight line into four equidistant segments from Point 285, Anterior Alar Point to Point 429, Posterior Symphyscal Point.

Anterior Inlet Point 456-458: The anterior most point subdividing a straight line into four equidistant segments from laint 285, Anterior Alar Point, to Point 429, Posterior Symphyseal Foint.

Inferior Isohial Inner Surface Point 459-461: The inferior most point of three equidistant points dividing a straight line between Point 396, Medial Tuberosity Point, and Point 450, Posterior Inlet Foint, into four equal segments.

Intermediate Ischial Inner Surface Point 462-464: The intermediate most point of three equidistant points dividing a straight line between Point 396, Medial Tuberosity Point, and Point 450, Posterior Inlet Point, into four equal segments.

Superior Ischial Inner Surface Point 365-567: The superior most point of three equidistant points dividing a straight line between Point 396, Medial Tuberosity Point, and Point 450, Posterior Inlet Point, into four equal segments.

Inferior Vertical Iliac Fossa Centour Point 468-470: The inferior most point of three equidistant points dividing a straight line between Point 339, Iliooristale Summum, to Point 150, Posterier Inlet Point, into four equal segments.

Intermediate Vertical Iliac Fossa Contour Foint 471-473: The intermediate most point of three equidistant points dividing a straight line from Point 339, Iliocistale Summum, to Foint 450, Fosterier Inlet Foint, into four equal segments.

Superior Vertical Iliae Fessa Contour Foint 474-476: The superior most point of three equidistant points dividing a straight line from Point 339, Iliocristale Summum, to Foint 450, Fosterior Inlet Foint, into four equal segments.

Posterior Transverse Iliac Fossa Contour Loint 477-479: The posterior most point of three equidistant points dividing a straight line from Point 210, Lateral Alar-Auricular Point, to Point 366, Posterior Superior Iliospinale, into four equal segments.

Intermediate Transverse Iliac Fossa Contour Point 480-482: The intermediate most point of three equidistant points dividing a straight line from Point 210, Lateral Alar-Auricular Point, to Point 366, Fosterior Superior Iliospinale, into four equal segments.

Anterior Transverse Iliac Fossa Contour Foint 493-485: The anterior most point of three equidistant points dividing a straight line from Point 210, Lateral Alar-Auricular Point, to Point 366, Posterior Superior Iliospinale, into four equal segments.

Anterior Tliac Notch Foint 486-488: The point of deepest indentation of the anterior iliac border between Foint 198, Right Iliospinale Summm, and Point 489, Anterior Inferior Iliospinale.

Iliospinale, Anterior Inferior - FF--101: This point is located at the apex of the anterior border of the anterior inferior iliac spine as viewed from the medial aspect of the innominate.

Anterior Iliac Base Point 402-004: This point is found at the apex of the angle formed by the inferior terminus of the suterior iliac margin and the superior surface of the acetabular margin at the base of the ilium.

Anterior Transverse Lateral Iliac Surface Contour Point 498-500: The anterior most point of three equidistant points dividing a straight line extending from Point 199, Right Iliospinale Surmur, to Point 366, Posterior Superior Iliospinale, into four equal segments.

Intermediate Transverse Lateral Iliac Surface Contour Point 501-503: The intermediate most point of three equidirtant points dividing a straight line extending from Point 199, Right Iliospinale Summam, to Point 366, Posterior Superior Iliospinale, into four equal segments.

Posterior Transverse Lateral Iliac Surface Contour Point 504-506: The posterior most point of three equidistant points dividing a straight line extending from Point 199, Right Iliespinale Surraum, to Point 366, Posterior Superior Iliespinale, into four equal segments.

Superior-Nost Vertical Lateral Ischio-Iliac Contour Point 507-509: The superior most point of four equidistant points dividing a straight line extending from Point 339, Iliocristale Summum, to Point 405, Superior Tuberosity Point, into five equal acgments.

Superior Intermediate Vertical Lateral Ischio-Iliac Contour Point 510-512: The intermediate superior point of four equidistant points dividing a straight line extending from Point 339, Iliccristale Surrum, to Point 405, Superior Tuberosity Point, into five equal segments.

Inferior Intermediate Vertical Lateral Ischio-Iliac Contour Point 513-515: The intermediate inferior point of four equidistant points dividing a straight line extending from Point 339, Iliocristale Summum, to Point 405, Superior Tuberosity Point, into five equal segments.

Inferior-Most Intermediate Vertical Lateral Ischio-Iliac Contour Point 516-518: The inferior most point of four equidistant points dividing a straight line extending from Point 339, Iliccristale Summum, to Point 405, Superior Tuberosity Point, into five equal segments.

Acetabulion, Arterier 519-521: The most anteriorly projecting point defined on the pulic portion of the acetabular rim. It is located by rotating the innominate bone on the esteemetric board from position 1 to position 2.

Acetabulion, Superior 522-524: The calibration hemisphere is positioned into the acetabulum so that the superior extremity of one of the diameter lines is opposite Point 519, Acetabulion Anterior. The point is marked on the acetabular rim closest to the superior diameter line when the innominate is oriented in the anatomical position.

Acetabulion, Fosterior 525-527: The calibration hemisphere is positioned into the acetabulum so that the anterior extremity of one of the diameter lines is opposite Point 519, Acetabulion Anterior. The point is marked on the acetabular rim closest to the posterior diameter line when the innominate is oriented in the anatomical position.

Acetabulion, Inferior 528-530: The calibration hemisphere is positioned into the acetabulum so that the anterior extremity of one of the diameter lines is opposite Point 519, Acetalulion Anterior. The point is marked on the acetabular rim closest to the inferior diameter lone when the innominate is oriented in the anatomical position.

Acetabulion, Center Point 531-533: The calibration hemisphere is positioned into the acetabulum so that the anterior extremity of one of the hemisphere lines is opposite Point 519, Acetabulion Anterior. Insert a marker through the H-point hole and mark the contact point on the interior surface of the acetabulum.

H-Point 534-536: A best-fit plaxiglass hemisphere is placed into the acetabulum of the right innominate bone and positioned so that the anterior extremity of one of the perpendicular diameter lines is opposite Point 519, Acetabulum Anterior. H-Point is the center point of the hemisphere surface.




Figure 8. Epoxy models of the small female, medium male, and large male pelves. Lateral View.




Figure 7. Epoxy models of the small female, medium male, and large male pelves. Anterior View.




Figure 9. Epoxy models of the small female, medium male, and large male pelves. Superior Niew.




Figure 9. Epoxy models of the small female, medium male, and large male pelves. Superior Niew.